

Trends in body mass index changes among Korean adolescents between 2005-2020, including the COVID-19 pandemic period: a national representative survey of one million adolescents

M.J. KIM¹, K.H. LEE², J.S. LEE³, N. KIM⁴, J.Y. SONG⁵, Y.H. SHIN⁵, J.M. YANG^{6,7}, S.W. LEE^{8,9}, J. HWANG¹⁰, S.Y. RHEE^{11,12}, D.K. YON^{8,12}, J.I. SHIN¹, Y.J. CHOI²

¹Department of Pediatrics, Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea

²Department of Pediatrics, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, South Korea

³Department of Pediatrics, Korea University Anam Hospital, Korea University College of Medicine, Seoul, South Korea

⁴Department of Neuropsychiatry, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, South Korea

⁵Department of Pediatrics, CHA Gangnam Medical Center, CHA University School of Medicine, Seoul, South Korea

⁶Department of Ophthalmology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, South Korea

⁷Department of Ophthalmology, Dongguk University Ilsan Hospital, Goyang, South Korea

⁸Department of Data Science, Sejong University College of Software Convergence, Seoul, South Korea

⁹Department of Precision Medicine, Sungkyunkwan University School of Medicine, Suwon, South Korea

¹⁰Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

¹¹Department of Endocrinology and Metabolism, Kyung Hee University School of Medicine, Seoul, South Korea

¹²Center for Digital Health, Medical Science Research Institute, Kyung Hee University Medical Center, Kyung Hee University College of Medicine, Seoul, South Korea

Abstract. – **OBJECTIVE:** The impact of the coronavirus disease 2019 (COVID-19) pandemic on weight gain in children and adolescents remains unknown. We aimed to identify an estimated 15-year trend in mean body mass index (BMI) changes and prevalence of obesity and overweight among Korean adolescents from 2005 to 2020, including the period of the COVID-19 pandemic.

PATIENTS AND METHODS: We analyzed data taken from a nationwide survey (Korea Youth Risk Behavior Survey), between 2005 and 2020. Representative samples of one million Korean adolescents aged 13-18 years (n=1,057,885) were examined. The 15-year trends in mean BMI and proportion of obesity or overweight, and the changes due to the COVID-19 pandemic were analyzed.

RESULTS: The data of 1,057,885 Korean adolescents were analyzed (mean age: 14.98 years; females, 48.4%). The estimated weighted mean BMI was 20.5 kg/m² [95% confidence interval (CI), 20.4-20.5] from 2005 to 2008 and 21.5 kg/m² (95% CI,

21.4-21.6) in 2020 (during the COVID-19 pandemic). Although the 15-year trend of mean BMI gradually increased, the change in mean BMI before and during the pandemic significantly lessened (β diff, -0.027; 95% CI, -0.028 to -0.026). The 15-year (2005-2020) trend changes in the prevalence of obesity and overweight were similar (obesity prevalence from 2005-2008, 3.2%; 95% CI, 3.1-3.3 vs. obesity prevalence in 2020, 8.6%; 95% CI, 8.2-9.0; β diff, -0.309; 95% CI, -0.330 to -0.288).

CONCLUSIONS: The 15-year trend of overall mean BMI and obesity and overweight prevalence demonstrated a significant increase; however, its slope decreased during the pandemic. These landmark results suggest the need for the development of precise strategies to prevent pediatric obesity and overweight during the COVID-19 pandemic.

Key Words:

COVID-19, BMI, Obesity, Overweight, Adolescent, KYRBS.

Corresponding Authors: Dong Keon Yon, MD, FACAAl; e-mail: yonkkang@gmail.com
Jae Il Shin, MD, Ph.D; e-mail: SHINJI@yuhs.ac
Yun Jung Choi, MD; e-mail: flubber224@gmail.com

Introduction

The prevalence of obesity and overweight in children and adolescents is increasing worldwide, and the public health consequences are becoming a major issue globally^{1,2}. Obesity and overweight are influenced by complex interactions among diverse factors, such as lifestyle, physical activity, and comorbidities³. Obese children, especially adolescents, are likely to remain obese adults, and the adverse physical and psychological effects related to obesity are likely to persist into adulthood⁴. Therefore, it is important to identify the rate of obesity in adolescents and childhood for preventing adulthood obesity.

The coronavirus disease 2019 (COVID-19) pandemic has drastically altered the behavioral traits of people across society^{3,5}. In line with government policies for social distancing, changes such as school closures, disrupted dietary habits, and decreased physical activity increased the risk of weight gain⁶. Moreover, adolescents followed an alternate schedule for online classes and cut back on all activities outside their homes, including social gatherings⁷. Several scholars⁸ have reported the association of COVID-19 with weight gain in adults. Although adolescents are expected to gain more weight due to restricted activities, such as online classes and social distancing, studies on this age group are lacking⁹.

Therefore, this study aimed to investigate an estimated 15-year trend in mean body mass index (BMI) changes and the prevalence of obesity and overweight among one million Korean adolescents. We used data from the national representative Korea Youth Risk Behavior Survey (KYRBS) disseminated from 2005 to 2020, including the COVID-19 pandemic period. Additionally, we evaluated the pandemic-related weight changes in adolescents and compared the BMI trends of Korean adolescents during the pre-pandemic period (2005-2009) and during the pandemic (2020).

Patients and Methods

Study Population

We used the 2005-2020 data from the KYRBS¹⁰, supervised by the Korea Disease Control and Prevention Agency (KDCA) and Ministry of Education, to establish and evaluate government policies. Data were used to investigate health behaviors of a national representative sample of adolescents. The KYRBS selected the study pop-

ulation by applying a two-step stratification in the order of school and class, with details specified on the official website of the KDCA (<http://www.kdca.go.kr/yhs/>). The participants completed a web-based survey in computer laboratories at their respective schools. The KYRBS sampling procedures were performed to account for sample weights, clustering, and stratification to provide national representative estimates of the total Korean adolescent population¹⁰. These data were voluntarily gathered with a response rate of over 95%¹⁰. All adolescents (age 12-18 years, middle-high school) in this database were included in the present study. Participants with missing data were excluded from the study. The study protocol was approved by the KCDA and Sejong University (SJU-HR-E-2020-003).

Variable Definitions

Body mass index (BMI) was calculated from the participant's self-reported height and weight and classified, using age- and sex-specific equations from the 2007 Korean national growth charts, into: underweight (< 5th percentile), normal (5-84th percentile), overweight (85-94th percentile), and obesity (\geq 95th percentile)¹¹. The areas of residence were classified as urban (Seoul, Gyeonggi, Busan, Daegu, Incheon, Gwangju, Daejeon, Ulsan, and Sejong) or rural (Gangwon, Chungbuk, Chungnam, Jeonbuk, Jeonnam, Gyeongbuk, Gyeongnam, and Jeju) areas^{12,13}. Smoking was defined as having smoked a cigarette at least once within the last 30 days¹⁴. Parents' educational levels were determined based on the highest educational level of either parent (high school or lower, college or higher, and unknown)¹⁰. Economic status and school performance were classified as high, middle-high, middle, middle-low, or low. Parents' educational levels, economic status, and school performance were self-reported¹⁵.

Statistical Analysis

We used KYRBS data (2005-2020) to calculate the trend of mean BMI or proportion of obesity and overweight, stratified by sex, grade, and area of residence. The primary endpoint was the change in the mean BMI over a 15-year period, and the secondary endpoints were the 15-year trends for prevalence of obesity and overweight. We aimed to evaluate the impact of COVID-19 pandemic on the trends of obesity prevalence and compared these trends before and during the pandemic.

To stabilize the prevalence of obesity and overweight, the pre-COVID-19 period was set based on the KYRBS cycle of four consecutive years. Weighted complex sampling analysis with binary and linear logistic regression models and Fisher's exact tests were performed. The results are presented as weighted odds ratios (ORs) with 95% confidence intervals (CIs) or weighted β -coefficients with 95% CIs. In linear regression, the KYRBS cycle was included as a continuous variable [2005-2008, 2009-2012, 2013-2016, 2017-2019, and 2020 (COVID-19 pandemic)]. For binary regression, the KYRBS cycle was included (2017-2019 vs. COVID-19 pandemic). All the analyses were performed using SPSS version 25.0 (IBM Corp., Armonk, NY, USA) and R software V.3.1.1 (R Foundation, Vienna, Austria)¹⁶⁻¹⁸. Statistical significance was defined as a two-sided p -value of <0.05 .

Patient and Public Involvement

None of the patients were directly involved in designing the research question or conducting the research. Patients were not asked for advice on interpretation or writing of the results. There were no plans to involve patients or the relevant patient community in the dissemination of study findings at this time.

Results

From 2005 to 2020, a total of 1,057,885 adolescents were enrolled in the KYRBS, comprising 48.4% females, with a mean age of 14.98 years [standard deviation (SD): 1.74]. There were 545,460 (51.6%) participants in the 7-9th grade (middle school) and 512,425 (48.4%) in the 10-12th grade (high school). By area of residence, 573,274 (54.2%) participants lived in rural areas and 484,611 (45.8%) lived in urban areas. Based on their BMI, 53,197 (5.0%) participants were underweight ($<5^{\text{th}}$ percentile), 854,315 (80.8%) had a normal weight (5-85th percentile), and 101,930 (9.6%) were overweight (85-95th percentile). Table I presents the general characteristics of the study population.

Table II illustrates the national crude mean and overall, 15-year trend of the mean BMI in Korean adolescents from 2005 to 2020, including the COVID-19 pandemic period. The estimated national weighted mean BMI was 20.5 (95% CI, 20.4-20.5) between 2005 and 2008, and 21.5 (95% CI, 21.4-21.6) in 2020. Although the 15-year trend

of the mean BMI showed a gradual increase over time, the change in mean BMI before and during the pandemic was significantly decreased (β diff, -0.027; 95% CI, -0.028 to -0.026). The difference in the slope was consistent regardless of sex (male: β diff, -0.022; 95% CI, -0.025 to -0.019; female: β diff, -0.037; 95% CI, -0.040 to -0.035), grade (7-9th grade: β diff, -0.015; 95% CI, -0.017 to -0.013; 10-12th grade: β diff, -0.032; 95% CI, -0.034 to -0.030), and area of residence (rural: β diff, -0.031; 95% CI, -0.034 to -0.029; urban: β diff, -0.022; 95% CI, -0.025 to -0.020).

The changes in the prevalence of obesity and overweight between 2005 and 2020 are shown in Tables III and IV and Figure 1. The national weighted prevalence of obesity was 3.2% (95% CI, 3.1-3.3) between 2005 and 2008 and 8.6% (95% CI, 8.2-9.0) in 2020. The proportion of participants with obesity increased from 2005 to 2020; however, the slope of the increasing trend was significantly decreased before and during the pandemic (β diff, -0.309; 95% CI, -0.330 to -0.288). This trend was consistent regardless of sex, (male: β diff, -0.265; 95% CI, -0.284 to -0.246; female: β diff, -0.511; 95% CI, -0.544 to -0.478), grade (7-9th grade: β diff, -0.241; 95% CI, -0.272 to -0.210; 10-12th grade: β diff, -0.316; 95% CI, -0.368 to -0.310), and area of residence (rural: β diff, -0.339; 95% CI, -0.368 to -0.310; urban: β diff, -0.273; 95% CI, -0.304 to -0.242).

The national weighted prevalence of overweight and obesity was 11.8% (95% CI, 11.6-12.0) between 2005 and 2008 and 21.4% (95% CI, 20.8-22.1) in 2020. The slope for the prevalence of overweight and obesity had been positive for 15 years, but the prevalence of overweight and obesity between pre-pandemic and during the pandemic had significantly decreased (β diff, -0.180; 95% CI, -0.194 to -0.166).

Discussion

Although several small studies on the short-term trend of prevalence of pediatric obesity have been reported⁹, this is the first large-scale national representative study to compare the long-term trends in BMI changes and the prevalence of obesity and overweight among adolescents during the COVID-19 pandemic period. In this study, the 15-year trends in overall mean BMI demonstrated a significant increase, from 20.5 in 2005-2008 to 21.5 in 2020, across different sexes, age groups, and areas of residence. However, the slope of the

Table 1. General characteristics of Korean adolescent participants in the KYRBS from 2005 to 2020 (total sample size = 1,057,885).

General characteristic	Total crude sample, n (%)
Total number, n (%)	1,057,885 (100.0)
Age, years, crude mean (SD)	14.98 (1.74)
Grade, crude number (%)	
7-9 th grade (middle school)	545,460 (51.6)
10-12 th grade (high school)	512,425 (48.4)
Sex, crude number (%)	
Female	512,520 (48.4)
Male	545,365 (51.6)
BMI, kg/m ² ; crude number (%)	
Underweight (below 5 th percentile)	53,197 (5.0)
Normal (5-85 th percentile)	854,315 (80.8)
Overweight (85-95 th percentile)	101,930 (9.6)
Obese (above 95 th percentile)	48,434 (4.6)
Region of residence, crude number (%)	
Rural	573,274 (54.2)
Urban	484,611 (45.8)
Smoking, crude number (%)	228,273 (21.6)
Highest educational level of parents, crude number (%)	
High school or lower	381,258 (36.0)
College or higher	516,598 (48.8)
Unknown	160,029 (15.1)
Economic level, crude number (%)	
High	82,133 (7.8)
Middle-high	276,907 (26.2)
Middle	476,776 (45.1)
Middle-low	175,335 (16.6)
Low	46,734 (4.4)
School performance, crude number (%)	
High	129,114 (12.2)
Middle-high	268,672 (25.4)
Middle	300,760 (28.4)
Middle-low	248,238 (23.5)
Low	111,101 (10.5)

BMI, body mass index; KYRBS, Korea Youth Risk Behavior Web-based Survey; SD, standard deviation.

increase in mean BMI and prevalence of obesity and overweight decreased during the COVID-19 pandemic.

BMI is a simple and widely used method for estimating body fat mass in men and women of any age. In the Asia-Pacific region, adult obesity is defined as a BMI of 25 or more, and overweight is defined as a BMI of 23-24.^{9,19} However, considering the continuous increase in height and weight in the pediatric population, obesity is assessed using percentiles of BMI by sex and age¹¹. For adolescents, a BMI that is above the 95th percentile is considered obese, under the 5th percentile is underweight, and between 85-94.5 is overweight¹¹. Our study analyzed 15-years' data, including the COVID-19 pandemic period of 2020, thereby allowing perception of not only the long-term trends but also the impact of the

pandemic on BMI changes. In a recent study⁹ on BMI changes based on pediatric population-based Kaiser Permanente Southern California electronic health record data, the COVID-19 pandemic was associated with a significant weight gain in the pediatric population, especially among the youngest children. These results are similar to the results of our study, though we found that the slope was significantly decreased during the pandemic.

In Korea, several studies on the prevalence of pediatric obesity have been reported²⁰. Although one study²¹ reported that pediatric obesity was on the rise until the early 2000s¹⁸, another showed that the prevalence of overweight and obesity among children and adolescents plateaued between 2001 and 2014²². Consistent with our study, another previous global study²³ demonstrated a

Table II. National weighted trend of mean BMI (95% CI) in the Korean adolescent population through the KYRBS between 2005 and 2020.

	Trends in mean BMI								
	2005 to 2008	2009 to 2012	2013 to 2016	2017 to 2019	2020 (COVID-19 pandemic)	Pre-pandemic trend, β (95% CI) ^a	Pandemic trend, β (95% CI) ^a	Trend difference, β_{diff} (95% CI)	2017 to 2019 vs. 2020 (COVID-19 pandemic), OR (95% CI) ^b
BMI, kg/m ² ; weighted mean (95% CI)									
Overall	20.5 (20.4 to 20.5)	20.5 (20.4 to 20.5)	20.8 (20.8 to 20.9)	21.3 (21.2 to 21.3)	21.5 (21.4 to 21.6)	0.030 (0.029 to 0.031)	0.003 (0.002 to 0.004)	-0.027 (-0.028 to -0.026)	1.016 (1.010 to 1.022)
Sex									
Male	20.8 (20.8 to 20.9)	20.9 (20.8 to 20.9)	21.2 (21.1 to 21.2)	21.8 (21.8 to 21.9)	22.3 (22.2 to 22.3)	0.028 (0.027 to 0.030)	0.006 (0.005 to 0.007)	-0.022 (-0.025 to -0.019)	1.031 (1.026 to 1.036)
Female	20.0 (20.0 to 20.1)	20.1 (20.0 to 20.1)	20.5 (20.4 to 20.5)	20.7 (20.7 to 20.8)	20.6 (20.6 to 20.7)	0.035 (0.033 to 0.037)	-0.002 (-0.003 to 0.000)	-0.037 (-0.040 to -0.035)	0.991 (0.984 to 0.999)
Grade, weighted % (95% CI)									
7 th -9 th grade (middle school)	20.0 (19.9 to 20.0)	20.0 (19.9 to 20.0)	20.2 (20.1 to 20.2)	20.6 (20.5 to 20.6)	21.0 (20.9 to 21.0)	0.021 (0.019 to 0.022)	0.006 (0.005 to 0.007)	-0.015 (-0.017 to -0.013)	1.032 (1.025 to 1.039)
10 th -12 th grade (high school)	21.1 (21.0 to 21.1)	21.0 (21.0 to 21.1)	21.4 (21.4 to 21.5)	21.9 (21.9 to 22.0)	22.0 (22.0 to 22.1)	0.034 (0.032 to 0.035)	0.002 (0.001 to 0.003)	-0.032 (-0.034 to -0.030)	1.009 (1.000 to 1.017)
Region of residence									
Rural	20.4 (20.3 to 20.4)	20.5 (20.5 to 20.5)	20.8 (20.8 to 20.9)	21.3 (21.3 to 21.4)	21.5 (21.4 to 21.5)	0.033 (0.031 to 0.035)	0.002 (0.001 to 0.004)	-0.031 (-0.034 to -0.029)	1.013 (1.006 to 1.021)
Urban	20.5 (20.5 to 20.6)	20.5 (20.5 to 20.6)	20.9 (20.8 to 20.9)	21.3 (21.2 to 21.3)	21.5 (21.4 to 21.6)	0.026 (0.024 to 0.028)	0.004 (0.002 to 0.005)	-0.022 (-0.025 to -0.020)	1.020 (1.011 to 1.029)

BMI, body mass index; CI, confidence interval; KYRBS, Korea Youth Risk Behavior Web-based Survey; OR, odds ratio. ^aEstimated β (95% CI) was derived using linear regression, and this analysis included the KYRBS cycle (2005-2008, 2009-2012, 2013-2016, 2017-2019, and 2020) as a continuous variable. ^bEstimated OR (95% CI) was calculated using binomial regression, and this analysis included the KYRBS cycle (2017-2019 vs. 2020) as a categorical variable. Numbers in bold indicate a significant difference ($p < 0.05$).

Trends in BMI changes among Korean children and adolescents

Table III. National weighted prevalence and trend of obesity (95% CI) in the Korean adolescent population through the KYRBS between 2005 and 2020.

	Trends in obesity								
	2005 to 2008	2009 to 2012	2013 to 2016	2017 to 2019	2020 (COVID-19 pandemic)	Pre-pandemic trend, β (95% CI) ^a	Pandemic trend, β (95% CI) ^a	Trend difference, β_{diff} (95% CI)	2017 to 2019 vs. 2020 (COVID-19 pandemic), OR (95% CI) ^b
Obesity, weighted % (95% CI)									
Overall	3.2 (3.1 to 3.3)	3.3 (3.2 to 3.4)	4.7 (4.6 to 4.9)	7.3 (7.1 to 7.5)	8.6 (8.2 to 9.0)	0.343 (0.325 to 0.361)	0.034 (0.023 to 0.046)	-0.309 (-0.330 to -0.288)	1.200 (1.132 to 1.272)
Sex									
Male	5.1 (5.0 to 5.3)	5.1 (5.0 to 5.3)	7.0 (6.8 to 7.2)	10.6 (10.4 to 10.9)	12.7 (12.2 to 13.2)	0.303 (0.287 to 0.320)	0.038 (0.028 to 0.048)	-0.265 (-0.284 to -0.246)	1.225 (1.163 to 1.291)
Female	1.0 (0.9 to 1.1)	1.2 (1.1 to 1.2)	2.3 (2.2 to 2.4)	3.7 (3.5 to 3.8)	4.2 (3.9 to 4.5)	0.537 (0.509 to 0.566)	0.026 (0.009 to 0.044)	-0.511 (-0.544 to -0.478)	1.151 (1.052 to 1.260)
Grade, weighted % (95% CI)									
7-9 th grade (middle school)	2.4 (2.3 to 2.5)	2.6 (2.5 to 2.7)	3.4 (3.3 to 3.5)	5.2 (5.0 to 5.4)	6.8 (6.4 to 7.3)	0.300 (0.275 to 0.326)	0.059 (0.041 to 0.077)	-0.241 (-0.272 to -0.210)	1.348 (1.237 to 1.468)
10-12 th grade (high school)	4.1 (3.9 to 4.4)	3.9 (3.7 to 4.1)	5.9 (5.7 to 6.2)	9.1 (8.8 to 9.4)	10.4 (9.8 to 11.0)	0.341 (0.317 to 0.365)	0.025 (0.011 to 0.039)	-0.316 (-0.344 to -0.288)	1.151 (1.067 to 1.242)
Region of residence									
Rural	3.0 (2.9 to 3.2)	3.2 (3.1 to 3.3)	4.8 (4.6 to 4.9)	7.3 (7.0 to 7.6)	8.4 (7.9 to 8.9)	0.367 (0.342 to 0.391)	0.028 (0.013 to 0.043)	-0.339 (-0.368 to -0.310)	1.163 (1.075 to 1.258)
Urban	3.4 (3.2 to 3.5)	3.3 (3.2 to 3.5)	4.7 (4.5 to 4.9)	7.2 (6.9 to 7.5)	8.9 (8.3 to 9.5)	0.315 (0.288 to 0.341)	0.042 (0.025 to 0.059)	-0.273 (-0.304 to -0.242)	1.252 (1.147 to 1.366)

BMI, body mass index; CI, confidence interval; KYRBS, Korea Youth Risk Behavior Web-based Survey; OR, odds ratio. ^aEstimated β (95% CI) was derived using linear regression, and this analysis included the KYRBS cycle (2005-2008, 2009-2012, 2013-2016, 2017-2019, and 2020) as a continuous variable. ^bEstimated OR (95% CI) was calculated using binomial regression, and this analysis included the KYRBS cycle (2017-2019 vs. 2020) as a categorical variable. Numbers in bold indicate a significant difference ($p < 0.05$).

Table IV. National weighted prevalence and trend of overweight or obesity (95% CI) in the Korean adolescent population through the KYRBS between 2005 and 2020.

	Trends in overweight or obesity								2017 to 2019 vs. 2020 (COVID-19 pandemic), OR (95% CI) ^b
	2005 to 2008	2009 to 2012	2013 to 2016	2017 to 2019	2020 (COVID-19 pandemic)	Pre-pandemic trend, β (95% CI) ^a	Pandemic trend, β (95% CI) ^a	Trend difference, β_{diff} (95% CI)	
Overweight or obesity, weighted % (95% CI) Overall	11.8 (11.6 to 12.0)	11.9 (11.7 to 12.1)	14.6 (14.4 to 14.8)	18.9 (18.6 to 19.2)	21.4 (20.8 to 22.1)	0.209 (0.198 to 0.220)	0.029 (0.021 to 0.037)	-0.180 (-0.194 to -0.166)	1.171 (1.123 to 1.222)
Sex									
Male	15.7 (15.4 to 15.9)	15.8 (15.6 to 16.1)	18.8 (18.5 to 19.0)	24.7 (24.3 to 25.0)	29.0 (28.3 to 29.7)	0.204 (0.194 to 0.215)	0.041 (0.033 to 0.048)	-0.163 (-0.176 to -0.150)	1.248 (1.201 to 1.296)
Female	7.4 (7.2 to 7.6)	7.4 (7.2 to 7.6)	10.1 (9.9 to 10.3)	12.6 (12.3 to 13.0)	13.3 (12.7 to 13.8)	0.237 (0.221 to 0.253)	0.010 (0.000 to 0.020)	-0.227 (-0.246 to -0.208)	1.057 (1.001 to 1.118)
Grade, weighted % (95% CI)									
7-9 th grade (middle school)	9.8 (9.5 to 10.0)	10.0 (9.7 to 10.2)	11.5 (11.2 to 11.7)	14.5 (14.2 to 14.9)	18.3 (17.5 to 19.1)	0.162 (0.147 to 0.176)	0.053 (0.042 to 0.065)	-0.109 (-0.128 to -0.090)	1.315 (1.240 to 1.394)
10-12 th grade (high school)	14.3 (13.9 to 14.6)	13.8 (13.4 to 14.1)	17.5 (17.1 to 17.8)	22.7 (22.2 to 23.2)	24.5 (23.6 to 25.5)	0.219 (0.204 to 0.234)	0.018 (0.008 to 0.029)	-0.201 (-0.219 to -0.183)	1.108 (1.046 to 1.174)
Region of residence									
Rural	11.3 (11.0 to 11.6)	11.7 (11.5 to 12.0)	14.6 (14.3 to 14.9)	19.0 (18.5 to 19.4)	21.2 (20.4 to 22.0)	0.229 (0.214 to 0.244)	0.025 (0.015 to 0.036)	-0.204 (-0.222 to -0.186)	1.147 (1.087 to 1.211)
Urban	12.3 (12.0 to 12.6)	12.0 (11.7 to 12.3)	14.7 (14.3 to 15.0)	18.8 (18.3 to 19.3)	21.8 (20.8 to 22.8)	0.185 (0.169 to 0.201)	0.034 (0.021 to 0.047)	-0.151 (-0.170 to -0.132)	1.204 (1.125 to 1.289)

BMI, body mass index; CI, confidence interval; KYRBS, Korea Youth Risk Behavior Web-based Survey; OR, odds ratio. ^aWe defined Period I as 2009-2011; Period II, 2012-2014; Period III, 2015-2017; Period IV, 2018-2019; and Period V, 2020. ^aEstimated β (95% CI) was derived using linear regression, and this analysis included the KYRBS cycle (2005-2008, 2009-2012, 2013-2016, 2017-2019, and 2020) as a continuous variable. ^bEstimated OR (95% CI) was calculated using binomial regression, and this analysis included the KYRBS cycle (2017-2019 vs. 2020) as a categorical variable. Numbers in bold indicate a significant difference ($p < 0.05$).




Figure 1. Nationwide trends and prevalence of obesity and overweight among one million Korean adolescents from 2005 to 2020.

plateauing of BMI and obesity trends in children and adolescents²⁰. According to our study, the trend of increasing prevalence was maintained during the pre-pandemic period and pandemic periods; however, the slope of this increase notably declined during the pandemic. In other words, Korean adolescents gained less weight during the pandemic period, which is in contrast with previous studies^{24,25} which showed a notable increase in children's weight gain during the COVID-19 pandemic. Thus, this is a distinctive and significant finding of our study, and to the best of our knowledge, this is the only study with this finding. In fact, previous studies found that preschool and school-aged children had greater pandemic-associated increases in BMI than adolescents did²⁶. In other words, adolescents, who were examined in this study, were less affected by pandemic-associated increases in BMI.

With urbanization, rapid socio-economic transition, decreased physical activity, and altered nutritional habits, BMI has substantially increased over time^{3,27,28}. During the COVID-19 pandemic, the considerable reduction in physical activity, as well as longer periods of sedentary behavior caused by the social distancing regulations, may have been the causes for weight gain²⁹. Dietary habit changes have been another factor contributing to increased body weight gain. Additionally, increased time spent indoors owing to social distancing norms, unhealthy eating patterns, including delivered food or fast food, have been more common among adolescents³⁰. However, these factors do not explain our finding of a less steep

slope of the prevalence of obesity or overweight during the pandemic period. Accordingly, further research is necessary to elucidate the causes of this inclination.

The greatest strength of this study is the analysis of trends based on the KYRBS data of one million Korean adolescents, which is the only database of BMI changes since the outbreak of the COVID-19 pandemic. Globally, researchers^{31,32} have been unable to investigate BMI or obesity, due to various measures implemented to combat the COVID-19 pandemic, such as lockdowns.^{28,29} This national representative survey was possible because the data for the survey were collected in a non-face-to-face manner prior to the pandemic. Data were collected from adolescents going through puberty through anonymous self-reported online questionnaires completed at school computer laboratories, due to the sensitivity of the survey content. Thus, the long-term trends observed in this study were not biased by differences in the survey methods.

The present study had some limitations. First, data were collected from the adolescent population of a single country (South Korea). Nonetheless, the study showed that the trend of BMI changed during the pandemic compared to before the pandemic. Additionally, the study included only one-year data of the pandemic period; however, 2020 was the year that witnessed the greatest impact of COVID-19, since restrictions such as social distancing were imposed during the first outbreak. Finally, this study was based on self-reported answers, and we did not conduct actual measurements or objective assessments. However, the reliability of the KYRBS data used in our study has been confirmed, and the kappa value for allergic diseases has been reported to be 0.74-0.80³³.

Conclusions

This is the first large-scale national representative study comparing the long-term trends in BMI and the prevalence of obesity and overweight among one million adolescents, including the period of the COVID-19 pandemic. The 15-year trends in overall mean BMI and proportion of obesity and overweight demonstrated a significant increase; however, the slope of the increase in BMI decreased during the COVID-19 pandemic, irrespective of sex, age, and area of residence. These landmark results provided an improved ep-

idemiological understanding of pediatric obesity during the COVID-19 pandemic and suggest the need for the development of precise strategies to prevent pediatric obesity and overweight during the COVID-19 pandemic.

Conflict of Interest

The Authors declare that they have no conflict of interests.

Funding

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (NRF-2021R111A2059735). The funders had no role in the study design, data collection, data analysis, data interpretation, or writing of the report.

References

- 1) Collaboration NRF. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017; 390: 2627-2642.
- 2) Kim SY. Nationwide COVID-19 vaccination coverage and COVID-19 incidence in South Korea, January 2022: a national official report. *Life Cycle* 2022; 2: e2.
- 3) Lee SW, Lee J, Moon SY, Jin HY, Yang JM, Ogi-no S, Song M, Hong SH, Abou Ghayda R, Kronbichler A, Koyanagi A, Jacob L, Dragioti E, Smith L, Giovannucci E, Lee IM, Lee DH, Lee KH, Shin YH, Kim SY, Kim MS, Won HH, Ekelund U, Shin JI, Yon DK. Physical activity and the risk of SARS-CoV-2 infection, severe COVID-19 illness and COVID-19 related mortality in South Korea: a nationwide cohort study. *Br J Sports Med* 2021. [bjsports-2021-104203](https://doi.org/10.1136/bjsports-2021-104203). doi: 10.1136/bjsports-2021-104203. Epub ahead of print.
- 4) Zheng M, Lamb KE, Grimes C, Laws R, Bolton K, Ong KK, Campbell K. Rapid weight gain during infancy and subsequent adiposity: a systematic review and meta-analysis of evidence. *Obes Rev* 2018; 19: 321-332.
- 5) Lee SW, Kim SY, Moon SY, Yang JM, Ha EK, Jee HM, Shin JI, Cho SH, Yon DK, Suh DI. Estimating COVID-19 Infection and Severity Risks in Patients with Chronic Rhinosinusitis: A Korean Nationwide Cohort Study. *J Allergy Clin Immunol Pract* 2021; 9: 2262-2271.e2262.
- 6) Lee SW, Yuh WT, Yang JM, Cho YS, Yoo IK, Koh HY, Marshall D, Oh D, Ha EK, Han MY, Yon DK. Nationwide Results of COVID-19 Contact Tracing in South Korea: Individual Participant Data From an Epidemiological Survey. *JMIR Med Inform* 2020; 8: e20992.
- 7) Han E, Tan MMJ, Turk E, Sridhar D, Leung GM, Shibuya K, Asgari N, Oh J, Garcia-Basteiro AL, Hanefeld J, Cook AR, Hsu LY, Teo YY, Heymann D, Clark H, McKee M, Legido-Quigley H. Lessons learnt from easing COVID-19 restrictions: an analysis of countries and regions in Asia Pacific and Europe. *Lancet* 2020; 396: 1525-1534.
- 8) Marchitelli S, Mazza C, Lenzi A, Ricci E, Gnessi L, Roma P. Weight Gain in a Sample of Patients Affected by Overweight/Obesity with and without a Psychiatric Diagnosis during the Covid-19 Lockdown. *Nutrients* 2020; 12.
- 9) Woolford SJ, Sidell M, Li X, Else V, Young DR, Resnicow K, Koebrnick C. Changes in Body Mass Index Among Children and Adolescents During the COVID-19 Pandemic. *JAMA* 2021; 326: 1434-1436.
- 10) Noh H, An J, Kim MJ, Sheen YH, Yoon J, Welsh B, Koh HY, Han MY, Yon DK. Sleep problems increase school accidents related to allergic diseases. *Pediatr Allergy Immunol* 2020; 31: 98-103.
- 11) Kim K, Yun SH, Jang MJ, Oh KW. Body fat percentile curves for Korean children and adolescents: a data from the Korea National Health and Nutrition Examination Survey 2009-2010. *J Korean Med Sci* 2013; 28: 443-449.
- 12) Yoo HW, Jin HY, Yon DK, Effenberger M, Shin YH, Kim SY, Yang JM, Kim MS, Koyanagi A, Jacob L, Smith L, Yoo IK, Shin JI, Lee SW. Non-alcoholic Fatty Liver Disease and COVID-19 Susceptibility and Outcomes: a Korean Nationwide Cohort. *J Korean Med Sci* 2021; 36: e291.
- 13) Yoo IK, Marshall DC, Cho JY, Yoo HW, Lee SW. N-Nitrosodimethylamine-contaminated ranitidine and risk of cancer in South Korea: a nationwide cohort study. *Life Cycle* 2021; 1: e1.
- 14) Shin YH, Shin JI, Moon SY, Jin HY, Kim SY, Yang JM, Cho SH, Kim S, Lee M, Park Y, Kim MS, Won HH, Hong SH, Kronbichler A, Koyanagi A, Jacob L, Smith L, Lee KH, Suh DI, Lee SW, Yon DK. Autoimmune inflammatory rheumatic diseases and COVID-19 outcomes in South Korea: a nationwide cohort study. *Lancet Rheumatol* 2021; 3: e698-e706.
- 15) Lee SW, Moon SY, Yon DK, Asthma, Disease R. Epidemiological and clinical characteristics of 5,628 patients with coronavirus disease 2019 in South Korea: A nationwide multicenter study. *Allergy, Asthma & Respiratory Disease* 2021; 9: 136-140.
- 16) Lee SW, Yang JM, Yoo IK, Moon SY, Ha EK, Yeniova A, Cho JY, Kim MS, Shin JI, Yon DK. Proton pump inhibitors and the risk of severe COVID-19: a post-hoc analysis from the Korean nationwide cohort. *Gut* 2021; 70: 2013-2015.
- 17) Lee SW, Yang JM, Moon SY, Kim N, Ahn YM, Kim JM, Shin JI, Suh DI, Yon DK. Association between mental illness and COVID-19 in South Korea: a post-hoc analysis. *Lancet Psychiatry* 2021; 8: 271-272.

- 18) Lee SW. Methods for testing statistical differences between groups in medical research: statistical standard and guideline of life cycle committee. *Life Cycle* 2022; 2: e1.
- 19) Woo A, Lee SW, Koh HY, Kim MA, Han MY, Yon DK. Incidence of cancer after asthma development: 2 independent population-based cohort studies. *J Allergy Clin Immunol* 2021; 147: 135-143.
- 20) Kim JH, Moon JS. Secular Trends in Pediatric Overweight and Obesity in Korea. *J Obes Metab Syndr* 2020; 29: 12-17.
- 21) Nam GE, Kim YH, Han K, Jung JH, Rhee EJ, Lee WY. Obesity Fact Sheet in Korea, 2020: Prevalence of Obesity by Obesity Class from 2009 to 2018. *J Obes Metab Syndr* 2021; 30: 141-148.
- 22) Nam HK, Kim HR, Rhie YJ, Lee KH. Trends in the prevalence of extreme obesity among Korean children and adolescents from 2001 to 2014. *J Pediatr Endocrinol Metab* 2017; 30: 517-523.
- 23) Chung A, Backholer K, Wong E, Palermo C, Keating C, Peeters A. Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: a systematic review. *Obes Rev* 2016; 17: 276-295.
- 24) Brooks CG, Spencer JR, Sprafka JM, Roehl KA, Ma J, Londhe AA, He F, Cheng A, Brown CA, Page J. Pediatric BMI changes during COVID-19 pandemic: An electronic health record-based retrospective cohort study. *EClinicalMedicine* 2021; 38: 101026.
- 25) Chang TH, Chen YC, Chen WY, Chen CY, Hsu WY, Chou Y, Chang YH. Weight Gain Associated with COVID-19 Lockdown in Children and Adolescents: A Systematic Review and Meta-Analysis. *Nutrients* 2021; 13.
- 26) Jenssen BP, Kelly MK, Powell M, Bouchelle Z, Mayne SL, Fiks AG. COVID-19 and Changes in Child Obesity. *Pediatrics* 2021; 147.
- 27) Lee SW, Kim SY, Moon SY, Yoo IK, Yoo EG, Eom GH, Kim JM, Shin JI, Jeong MH, Yang JM, Yon DK. Statin Use and COVID-19 Infectivity and Severity in South Korea: Two Population-Based Nationwide Cohort Studies. *JMIR Public Health Surveill* 2021; 7: e29379.
- 28) Choi MJ, Yang JW, Lee S, Kim JY, Oh JW, Lee J, Stubbs B, Lee KH, Koyanagi A, Hong SH, Ghayda RA, Hwang J, Dragioti E, Jacob L, Carvalho AF, Radua J, Thompson T, Smith L, Fornaro M, Stickley A, Bettac EL, Han YJ, Kronbichler A, Yon DK, Lee SW, Shin JI, Lee E, Solmi M. Suicide associated with COVID-19 infection: an immunological point of view. *Eur Rev Med Pharmacol Sci* 2021; 25: 6397-6407.
- 29) Yang S, Guo B, Ao L, Yang C, Zhang L, Zhou J, Jia P. Obesity and activity patterns before and during COVID-19 lockdown among youths in China. *Clin Obes* 2020; 10: e12416.
- 30) Lim S, Lim H, Després JP. Collateral Damage of the COVID-19 Pandemic on Nutritional Quality and Physical Activity: Perspective from South Korea. *Obesity (Silver Spring)* 2020; 28: 1788-1790.
- 31) Hwang J, Park SH, Lee SW, Lee SB, Lee MH, Jeong GH, Kim MS, Kim JY, Koyanagi A, Jacob L, Jung SY, Song J, Yon DK, Shin JI, Smith L. Predictors of mortality in thrombotic thrombocytopenia after adenoviral COVID-19 vaccination: the FAPIC score. *Eur Heart J* 2021; 42: 4053-4063.
- 32) Yoo HW, Shin JI, Yon DK, Lee SW. COVID-19 Morbidity and Severity in Patients With Nonalcoholic Fatty Liver Disease in South Korea: A Nationwide Cohort Study. *Clin Gastroenterol Hepatol* 2022; e1217-e1218.
- 33) Bae J, Joung H, Kim JY, Kwon KN, Kim YT, Park SW. Test-retest reliability of a questionnaire for the Korea Youth Risk Behavior Web-based Survey. *J Prev Med Public Health* 2010; 43: 403-410.